

On the category of comodules over corings

Robert Wisbauer

Abstract

It is well known that the category \mathcal{M}^C of right comodules over an A -coring \mathcal{C} , A an associative ring, is a subcategory of the category of left modules ${}^*\mathcal{C}\mathcal{M}$ over the dual ring ${}^*\mathcal{C}$. The main purpose of this note is to show that \mathcal{M}^C is a full subcategory in ${}^*\mathcal{C}\mathcal{M}$ if and only if \mathcal{C} is locally projective as a left A -module.

1 Introduction

For any coassociative coalgebra C over a commutative ring R , the convolution product turns the dual module $C^* = \text{Hom}_R(C, R)$ into an associative R -algebra. The category \mathcal{M}^C of right comodules is an additive subcategory of the category ${}_{C^*}\mathcal{M}$ of left C^* -modules. \mathcal{M}^C is an abelian (in fact a Grothendieck) category if and only if C is flat as an R -module. Moreover, \mathcal{M}^C coincides with ${}_{C^*}\mathcal{M}$ if and only if C is finitely generated and projective as an R -module (e.g. [11, Corollary 33]).

In case C is projective as an R -module, \mathcal{M}^C is a full subcategory of ${}_{C^*}\mathcal{M}$ and coincides with $\sigma[C^*C]$, the category of submodules of C -generated C^* -modules (e.g. [9, 3.15, 4.3]). It was well understood from examples that projectivity of C as an R -module was not necessary to achieve $\mathcal{M}^C = \sigma[C^*C]$ and that the equality holds provided C satisfies the α -condition, i.e., the canonical maps $N \otimes_R C \rightarrow \text{Hom}_{\mathbb{Z}}(C^*, N)$ are injective for all R -modules N (e.g. [1, Satz 2.2.13], [2, Section 2], [10, 3.2]). It will follow from our results that this condition is in fact equivalent to $\mathcal{M}^C = \sigma[C^*C]$ and also to C being *locally projective* as an R -module.

We do investigate the questions and results mentioned above in the more general case of comodules over any A -coring, A an associative ring, and it will turn out that the above observations remain valid almost literally in this extended setting.

2 Some module theory

Let A be any associative ring with unit and denote $(-)^* = \text{Hom}_A(-, A)$. We write ${}_A\mathcal{M}$ (\mathcal{M}_A) for the category of unital left (right) A -modules. I (or I_N) will denote the identity map (of the module N).

2.1. Canonical maps. For any left A -module K , consider the maps

$$\tilde{\varphi}_K : \begin{array}{ccccc} K & \xrightarrow{\varphi_K} & K^{**} & \xrightarrow{i} & A^{K^*}, \\ k & \mapsto & [f \mapsto f(k)] & \mapsto & (f(k))_{f \in K^*}. \end{array}$$

For any right A -module N define the maps

$$\begin{array}{ccc} \alpha_{N,K} : N \otimes_A K & \rightarrow \text{Hom}_{\mathbb{Z}}(K^*, N), & n \otimes k \mapsto [f \mapsto nf(k)], \\ \psi_N : N \otimes_A A^{K^*} & \rightarrow N^{K^*}, & n \otimes (a_f)_{f \in K^*} \mapsto (na_f)_{f \in K^*}. \end{array}$$

By the identification $\text{Map}(K^*, N) = N^{K^*}$ we have the commutative diagram

$$\begin{array}{ccc} N \otimes_A K & \xrightarrow{I \otimes \tilde{\varphi}_K} & N \otimes_A A^{K^*} & & n \otimes k & \longrightarrow & n \otimes (f(k))_{f \in K^*} \\ \alpha_{N,K} \downarrow & & \downarrow \psi_N & & \downarrow & & \downarrow \\ 0 \longrightarrow \text{Hom}_{\mathbb{Z}}(K^*, N) & \longrightarrow & N^{K^*}, & & [f \mapsto nf(k)] & \longrightarrow & (nf(k))_{f \in K^*}. \end{array}$$

2.2. Injectivity of $\alpha_{N,K}$. We stick to the notation above.

(1) The following are equivalent:

- (a) $\alpha_{N,K}$ is injective;
- (b) for $u \in N \otimes_A K$, $(I \otimes f)(u) = 0$ for all $f \in K^*$, implies $u = 0$.

(2) The following are equivalent:

- (a) For every finitely presented right A -module N , $\alpha_{N,K}$ is injective;
- (b) $\tilde{\varphi}_K : K \rightarrow A^{K^*}$ is a pure monomorphism.

Proof. (1) Let $u = \sum_{i=1}^r n_i \otimes k_i \in N \otimes_A K$. Then $(I \otimes f)(u) = \sum_{i=1}^r n_i f(k_i) = 0$, for all $f \in K^*$ if and only if $u \in \text{Ke } \alpha_{N,K}$.

(2) For N finitely presented, ψ_N is injective (bijective) and so $\alpha_{N,K}$ is injective if and only if $I_N \otimes \tilde{\varphi}_K$ is injective. Injectivity of $I_N \otimes \tilde{\varphi}_K$ for all finitely presented N characterizes $\tilde{\varphi}_K$ as a pure monomorphism (e.g., [8, 34.5]). \square

We say that K satisfies the α -condition provided $\alpha_{N,K}$ is injective for all right A -modules N . Such modules are named *universally torsionless (UTL)* in [4] and we recall some of their characterizations.

The module K is called *locally projective* (see [12]) if, for any diagram of left A -modules with exact lines

$$\begin{array}{ccccccc} 0 & \longrightarrow & F & \xrightarrow{i} & K & & \\ & & & & \downarrow g & & \\ & & L & \xrightarrow{f} & N & \longrightarrow & 0, \end{array}$$

where F is finitely generated, there exists $h : K \rightarrow L$ such that $g \circ i = f \circ h \circ i$.

Clearly every projective module is locally projective. From Garfinkel [4, Theorem 3.2] and Huisgen-Zimmermann [12, Theorem 2.1] we have the following characterizations of these modules which are also studied in Ohm-Bush [5] (as *trace modules*), and in Raynaud-Gruson [6] (as *modules plats et strictement de Mittag-Leffler*).

2.3. Locally projective modules. *For the left A -module K , the following are equivalent:*

- (a) K is locally projective;
- (b) K is a pure submodule of a locally projective module;
- (c) $\alpha_{N,K}$ is injective, for any right A -module N ;
- (d) $\alpha_{N,K}$ is injective, for any cyclic right A -module N ;
- (e) for each $m \in K$, we have $m \in K^*(m)K$;
- (f) for each finitely generated submodule $i : F \rightarrow K$, there exists $n \in \mathbb{N}$ and maps $\beta : R^n \rightarrow K$, $\gamma : K \rightarrow R^n$ with $\beta \circ \gamma \circ i = i$.

Recall the following observations. Notice that for a right noetherian ring A , every product of copies of A is locally projective as left A -module (e.g. [12, Corollary 4.3]).

2.4. Corollary. *Let K be a left A -module.*

- (1) *Every locally projective module is flat and a pure submodule of some product A^Λ , Λ some set.*
- (2) *If K is finitely generated, or A is left perfect, then K is locally projective if and only if K is projective.*

(3) For a right noetherian ring A , the following are equivalent:

- (a) K is locally projective;
- (b) K is a pure submodule of a product A^Λ , Λ some set.

The following facts from general category theory will be helpful (e.g., [7]). In any category \mathcal{A} , a morphism $f : A \rightarrow B$ is called a *monomorphism* if for any morphisms $g, h : C \rightarrow A$ the identity $f \circ g = f \circ h$ implies $g = h$.

In an additive category \mathcal{A} a morphism $\gamma : K \rightarrow A$ is called a *kernel* of $f : A \rightarrow B$ provided $f \circ \gamma = 0$ and, for every $g : C \rightarrow A$ with $f \circ g = 0$, there is exactly one $h : C \rightarrow K$ such that $g = \gamma \circ h$.

Recall the following well-known (and easily proved) observations.

2.5. Monomorphisms. Let \mathcal{A} be any category and $f : A \rightarrow B$ a morphism in \mathcal{A} . The following are equivalent:

- (a) f is a monomorphism;
- (b) the map $\text{Mor}(L, f) : \text{Mor}(L, A) \rightarrow \text{Mor}(L, B)$, $g \mapsto f \circ g$, is injective, for any $L \in \mathcal{A}$.

If \mathcal{A} is additive and has kernels, then (a)-(b) are equivalent to:

- (c) for the kernel $\gamma : K \rightarrow A$ of f , $K = 0$.

The basic properties of adjoint functors will be helpful.

2.6. Adjoint functors. Let \mathcal{A} and \mathcal{B} be any categories. Assume a functor $F : \mathcal{A} \rightarrow \mathcal{B}$ is right adjoint to a functor $G : \mathcal{B} \rightarrow \mathcal{A}$, i.e.,

$$\text{Mor}_{\mathcal{B}}(Y, F(X)) \simeq \text{Mor}_{\mathcal{A}}(G(Y), X), \text{ for any } X \in \mathcal{A}, Y \in \mathcal{B}.$$

Then

- (1) F preserves monomorphisms and products,
- (2) G preserves epimorphisms and coproducts.

For the study of comodules the following type of module categories is of particular interest.

2.7. The category $\sigma[K]$. For any left A -module K we denote by $\sigma[K]$ the full subcategory of ${}_A\mathcal{M}$ whose objects are submodules of K -generated modules. This is the smallest full Grothendieck subcategory of ${}_A\mathcal{M}$ containing K (see [8]).

$\sigma[K]$ coincides with ${}_A\mathcal{M}$ if and only if A embeds into some (finite) co-product of copies of K . This happens, for example, when K is a faithful A -module which is finitely generated as a module over its endomorphism ring (see [8, 15.4]).

The *trace functor* $\mathcal{T}^K : {}_A\mathcal{M} \rightarrow \sigma[K]$, which sends any $X \in {}_A\mathcal{M}$ to

$$\mathcal{T}^K(X) := \sum \{f(N) \mid N \in \sigma[K], f \in \text{Hom}_A(N, X)\},$$

is right adjoint to the inclusion functor $\sigma[K] \rightarrow {}_A\mathcal{M}$ (e.g., [8, 45.11]). Hence, by 2.6, for any family $\{N_\lambda\}_\Lambda$ of modules in $\sigma[K]$, the product in $\sigma[K]$ is

$$\prod_\Lambda^K N_\lambda = \mathcal{T}^K(\prod_\Lambda N_\lambda),$$

where the unadorned \prod denotes the usual (cartesian) product of A -modules.

It also follows from 2.6 that for $\{N_\lambda\}_\Lambda$ in $\sigma[K]$ the coproduct in $\sigma[K]$ and the coproduct in ${}_A\mathcal{M}$ coincide.

3 Corings and comodules

As before, let A be any associative ring with unit.

3.1. Corings and their duals. An *A-coring* is an (A, A) -bimodule \mathcal{C} with (A, A) -bimodule maps (*comultiplication* and *counit*)

$$\underline{\Delta} : \mathcal{C} \rightarrow \mathcal{C} \otimes_A \mathcal{C}, \quad \underline{\varepsilon} : \mathcal{C} \rightarrow A,$$

satisfying the identities

$$(I \otimes \underline{\Delta}) \circ \underline{\Delta} = (\underline{\Delta} \otimes I) \circ \underline{\Delta}, \quad (I \otimes \underline{\varepsilon}) \circ \underline{\Delta} = I = (\underline{\varepsilon} \otimes I) \circ \underline{\Delta}.$$

For elementwise description of these maps we adopt the Σ -notation, writing for $c \in \mathcal{C}$,

$$\underline{\Delta}(c) = \sum c_{\underline{1}} \otimes c_{\underline{2}}.$$

Then coassociativity of $\underline{\Delta}$ is written as

$$\sum \underline{\Delta}(c_{\underline{1}}) \otimes c_{\underline{2}} = \sum c_{\underline{1}\underline{1}} \otimes c_{\underline{1}\underline{2}} \otimes c_{\underline{2}} = \sum c_{\underline{1}} \otimes c_{\underline{2}\underline{1}} \otimes c_{\underline{2}\underline{2}} = \sum c_{\underline{1}} \otimes \underline{\Delta}(c_{\underline{2}}),$$

and the conditions on the counit are

$$\sum \underline{\varepsilon}(c_{\underline{1}}) c_{\underline{2}} = c = \sum c_{\underline{1}} \underline{\varepsilon}(c_{\underline{2}}).$$

Of course, when A is commutative and $ac = ca$ for all $a \in A$, $c \in \mathcal{C}$, the coring \mathcal{C} is just a *coalgebra* in the usual sense.

For any A -coring \mathcal{C} , the maps $\mathcal{C} \rightarrow A$ may be right A -linear or left A -linear and we denote these by

$$\mathcal{C}^* := \text{Hom}_{-A}(\mathcal{C}, A), \quad {}^*\mathcal{C} := \text{Hom}_{A-}(\mathcal{C}, A),$$

and for bilinear maps we have $\text{Hom}_{AA}(\mathcal{C}, A) = {}^*\mathcal{C} \cap \mathcal{C}^*$.

Both \mathcal{C}^* and ${}^*\mathcal{C}$ can be turned to associative rings with unit $\underline{\varepsilon}$ by the (convolution) products

- (1) for $f, g \in \mathcal{C}^*$, and $c \in \mathcal{C}$ put $f *^r g(c) = \sum g(f(c_1)c_2)$,
- (2) for $f, g \in {}^*\mathcal{C}$, and $c \in \mathcal{C}$ put $f *^l g(c) = \sum f(c_1g(c_2))$.

Notice that for $f, g \in {}^*\mathcal{C} \cap \mathcal{C}^*$ this yields

$$f * g(c) = \sum f(c_1)g(c_2),$$

a formula which is well known from coalgebras.

It is easily verified that the maps

$$\iota_l : A \rightarrow {}^*\mathcal{C}, \quad a \mapsto [c \mapsto \underline{\varepsilon}(c)a], \quad \text{and} \quad \iota_r : A \rightarrow \mathcal{C}^*, \quad a \mapsto [c \mapsto a\underline{\varepsilon}(c)],$$

are ring anti-morphisms and hence we may consider left ${}^*\mathcal{C}$ -modules as right A -modules and right \mathcal{C}^* -modules as left A -modules.

3.2. Right comodules. Let \mathcal{C} be an A -coring and M a right A -module. An A -linear map $\varrho_M : M \rightarrow M \otimes_A \mathcal{C}$ is called a *coaction* on M , and it is said to be *counital* and *coassociative* provided

$$(I \otimes \underline{\varepsilon}) \circ \varrho_M = I, \quad \text{and} \quad (I \otimes \underline{\Delta}) \circ \varrho_M = (\varrho_M \otimes I) \circ \varrho_M.$$

A *right \mathcal{C} -comodule* is a right A -module with a counital coassociative coaction.

A *morphism* of right \mathcal{C} -comodules $f : M \rightarrow N$ is an A -linear map such that

$$\varrho_N \circ f = (f \otimes I) \circ \varrho_M.$$

We denote the set of comodule morphisms between M and N by $\text{Hom}^{\mathcal{C}}(M, N)$. It is easy to show that this is an abelian group and hence the category $\mathcal{M}^{\mathcal{C}}$, formed by right \mathcal{C} -comodules and comodule morphisms, is additive.

For any right A -module X , the tensor product $X \otimes_A \mathcal{C}$ is a right \mathcal{C} -comodule by

$$I_{\otimes \underline{\Delta}} : X \otimes_A \mathcal{C} \rightarrow X \otimes_A \mathcal{C} \otimes_A \mathcal{C},$$

and for any A -morphism $f : X \rightarrow Y$, the map

$$f_{\otimes I} : X \otimes_A \mathcal{C} \rightarrow Y \otimes_A \mathcal{C}$$

is a comodule morphism.

3.3. The category $\mathcal{M}^{\mathcal{C}}$. Let \mathcal{C} be an A -coring.

(1) The category $\mathcal{M}^{\mathcal{C}}$ has direct sums and cokernels.

It has kernels provided \mathcal{C} is flat as a left A -module.

(2) For the functor $- \otimes_A \mathcal{C} : \mathcal{M}_A \rightarrow \mathcal{M}^{\mathcal{C}}$ we have natural isomorphisms

$$\mathrm{Hom}^{\mathcal{C}}(M, X \otimes_A \mathcal{C}) \rightarrow \mathrm{Hom}_A(M, X), \quad f \mapsto (I_{\otimes \underline{\Delta}}) \circ f,$$

for $M \in \mathcal{M}^{\mathcal{C}}$, $X \in \mathcal{M}_A$, with inverse map $h \mapsto (h_{\otimes I}) \circ \varrho_M$, i.e., the functor $- \otimes_A \mathcal{C} : \mathcal{M}_A \rightarrow \mathcal{M}^{\mathcal{C}}$ is right adjoint to the forgetful functor $\mathcal{M}^{\mathcal{C}} \rightarrow \mathcal{M}_A$ and hence it preserves monomorphisms and products.

(3) For the right comodule endomorphisms we have $\mathrm{End}^{\mathcal{C}}(\mathcal{C}) \simeq \mathcal{C}^*$.

(4) \mathcal{C} is a subgenerator in $\mathcal{M}^{\mathcal{C}}$.

Proof. (1) Consider a family $\{M_\lambda\}_\Lambda$ of right \mathcal{C} -comodules. It is easy to prove that the direct sum $\bigoplus_\Lambda M_\lambda$ in \mathcal{M}_A is a right \mathcal{C} -comodule and has the universal property of a coproduct in $\mathcal{M}^{\mathcal{C}}$.

For any morphism $f : M \rightarrow N$ of right \mathcal{C} -comodules, the cokernel of f in \mathcal{M}_A has a comodule structure and hence is a cokernel in $\mathcal{M}^{\mathcal{C}}$. If \mathcal{C} is flat as a left A -module, similar arguments hold for the kernel.

(2) The proof of the corresponding assertion for coalgebras applies (e.g., [9, 3.12]) and then refer to 2.6. Note that the adjointness, for example, was also observed in [3, Lemma 3.1].

(3) The group isomorphism $\mathrm{End}^{\mathcal{C}}(\mathcal{C}) \simeq \mathcal{C}^*$ follows from (2) by putting $M = \mathcal{C}$ and $X = A$. This is a ring isomorphism when writing the morphisms on the right.

(4) For any $M \in \mathcal{M}^{\mathcal{C}}$, there is an epimorphism $A^{(\Lambda)} \rightarrow M$ in \mathcal{M}_A . Tensoring with \mathcal{C} yields an epimorphism $A^{(\Lambda)} \otimes_A \mathcal{C} \rightarrow M \otimes_A \mathcal{C}$ in $\mathcal{M}^{\mathcal{C}}$. As easily checked the structure map $\varrho_M : M \rightarrow M \otimes_A \mathcal{C}$ is a morphism in $\mathcal{M}^{\mathcal{C}}$ and hence M is a subobject of a \mathcal{C} -generated comodule. \square

3.4. $\mathcal{M}^{\mathcal{C}}$ as Grothendieck category.

For an A -coring \mathcal{C} the following are equivalent:

- (a) \mathcal{C} is a flat left A -module;
- (b) every monomorphism in $\mathcal{M}^{\mathcal{C}}$ is injective;
- (c) the forgetful functor $\mathcal{M}^{\mathcal{C}} \rightarrow \mathcal{M}_A$ respects monomorphisms.

If these conditions are satisfied, $\mathcal{M}^{\mathcal{C}}$ is a Grothendieck category.

Proof. (a) \Rightarrow (b) \Leftrightarrow (c) are obvious.

(c) \Rightarrow (a) For any monomorphism $f : N \rightarrow L$ in \mathcal{M}_A , the map $f \otimes I : N \otimes_A \mathcal{C} \rightarrow L \otimes_A \mathcal{C}$ is a monomorphism in $\mathcal{M}^{\mathcal{C}}$ (by 3.3(2)) and hence injective by assumption. This shows that $- \otimes_A \mathcal{C} : \mathcal{M}_A \rightarrow \mathbb{Z}\text{-Mod}$ is exact and hence \mathcal{C} is a flat left A -module.

Now assume that (a)-(c) are satisfied. Then $\mathcal{M}^{\mathcal{C}}$ is abelian and cocomplete. Since \mathcal{C} is a subgenerator it is routine to show that the subcomodules of \mathcal{C}^n , $n \in \mathbb{N}$, form a generating set for $\mathcal{M}^{\mathcal{C}}$. Hence $\mathcal{M}^{\mathcal{C}}$ is a Grothendieck category. \square

Every right \mathcal{C} -comodule M allows a left ${}^*\mathcal{C}$ -module structure by

$$\rightarrow : {}^*\mathcal{C} \otimes_{\mathbb{Z}} M \rightarrow M, \quad f \otimes m \mapsto (I \otimes f) \circ \varrho_M(m).$$

With this structure any comodule morphisms $M \rightarrow N$ is ${}^*\mathcal{C}$ -linear, i.e.

$$\text{Hom}^{\mathcal{C}}(M, N) \subset \text{Hom}_{{}^*\mathcal{C}}(M, N),$$

and hence $\mathcal{M}^{\mathcal{C}}$ is a subcategory of ${}^*\mathcal{C}\mathcal{M}$. As shown in [3, Lemma 4.3], $\mathcal{M}^{\mathcal{C}}$ can be identified with ${}^*\mathcal{C}\mathcal{M}$ provided \mathcal{C} is finitely generated and projective as left A -module.

Notice that in any case \mathcal{C} is a faithful ${}^*\mathcal{C}$ -module since $f \rightarrow c = 0$ for all $c \in \mathcal{C}$ implies $f(c) = \underline{\varepsilon}(f \rightarrow c) = 0$ and hence $f = 0$.

The question arises when, more generally, $\mathcal{M}^{\mathcal{C}}$ is a full subcategory of ${}^*\mathcal{C}\mathcal{M}$, i.e., when $\text{Hom}^{\mathcal{C}}(M, N) = \text{Hom}_{{}^*\mathcal{C}}(M, N)$, for any $M, N \in \mathcal{M}^{\mathcal{C}}$. The answer is given in our main theorem:

3.5. $\mathcal{M}^{\mathcal{C}}$ as full subcategory of ${}^*\mathcal{C}\mathcal{M}$

For the A -coring \mathcal{C} , the following are equivalent:

- (a) $\mathcal{M}^{\mathcal{C}} = \sigma[{}^*\mathcal{C}\mathcal{C}]$;
- (b) $\mathcal{M}^{\mathcal{C}}$ is a full subcategory of ${}^*\mathcal{C}\mathcal{M}$;

- (c) for all $M, N \in \mathcal{M}^{\mathcal{C}}$, $\text{Hom}^{\mathcal{C}}(M, N) = \text{Hom}_{*\mathcal{C}}(M, N)$;
- (d) \mathcal{C} satisfies the α -condition as left A -module;
- (e) every $*\mathcal{C}$ -submodule of \mathcal{C}^n , $n \in \mathbb{N}$, is a subcomodule of \mathcal{C}^n ;
- (f) \mathcal{C} is locally projective as left A -module.

If these conditions are satisfied we have, for any family $\{M_\lambda\}_\Lambda$ of right A -modules,

$$\left(\prod_{\Lambda} M_\lambda\right) \otimes_A \mathcal{C} \simeq \prod_{\Lambda}^{\mathcal{C}} (M_\lambda \otimes_A \mathcal{C}) \subset \prod_{\Lambda} (M_\lambda \otimes_A \mathcal{C}).$$

Proof. The implications (a) \Leftrightarrow (b) \Leftrightarrow (c) \Rightarrow (e) are obvious.

(a) \Rightarrow (d) By 3.4 ${}_A\mathcal{C}$ is flat. For any $N \in \mathcal{M}_A$ we prove the injectivity of the map

$$\alpha : N \otimes_A \mathcal{C} \rightarrow \text{Hom}_{\mathbb{Z}}(*\mathcal{C}, N), \quad n \otimes c \mapsto [f \mapsto nf(c)].$$

Considering $\text{Hom}_{\mathbb{Z}}(*\mathcal{C}, N)$ and the right \mathcal{C} -comodule $N \otimes_A \mathcal{C}$ as left $*\mathcal{C}$ -modules in the canonical way, we observe that α is $*\mathcal{C}$ -linear. So for any right \mathcal{C} -comodule L we have the commutative diagram

$$\begin{array}{ccc} \text{Hom}_{*\mathcal{C}}(L, N \otimes_A \mathcal{C}) & \xrightarrow{\text{Hom}(L, \alpha)} & \text{Hom}_{*\mathcal{C}}(L, \text{Hom}_{\mathbb{Z}}(*\mathcal{C}, N)) \\ \downarrow \simeq & & \downarrow \simeq \\ \text{Hom}_A(L, N) & \xrightarrow{i} & \text{Hom}_{\mathbb{Z}}(L, N), \end{array}$$

where the first vertical isomorphism is obtained by assumption and 3.3,

$$\text{Hom}_{*\mathcal{C}}(L, N \otimes_A \mathcal{C}) = \text{Hom}^{\mathcal{C}}(L, N \otimes_A \mathcal{C}) \simeq \text{Hom}_A(L, N),$$

and the second one by canonical isomorphisms

$$\text{Hom}_{*\mathcal{C}}(L, \text{Hom}_{\mathbb{Z}}(*\mathcal{C}, N)) \simeq \text{Hom}_{\mathbb{Z}}(*\mathcal{C} \otimes_{*\mathcal{C}} L, N) \simeq \text{Hom}_{\mathbb{Z}}(L, N).$$

This shows that $\text{Hom}(L, \alpha)$ is injective and so (by 2.5) the corestriction of α is a monomorphism in $\mathcal{M}^{\mathcal{C}}$. Since ${}_A\mathcal{C}$ is flat this implies that α is injective (by 3.4).

(e) \Rightarrow (a) First we show that every finitely generated module $N \in \sigma[*\mathcal{C}]$ is a \mathcal{C} -comodule. There exists some $*\mathcal{C}$ -submodule $X \subset \mathcal{C}^n$, $n \in \mathbb{N}$, and

an epimorphism $h : X \rightarrow N$. By assumption X and the kernel of h are comodules and hence N is a comodule.

Now for any $L \in \sigma[*_{\mathcal{C}}\mathcal{C}]$ the finitely generated submodules are comodules and hence L is a comodule.

For any $*_{\mathcal{C}}$ -morphism in $\sigma[*_{\mathcal{C}}\mathcal{C}]$, the kernel is a $*_{\mathcal{C}}$ -submodule and hence a comodule. As easily verified this implies that monomorphisms and epimorphisms in $\sigma[*_{\mathcal{C}}\mathcal{C}]$ are comodule morphisms and hence this is true for all morphisms in $\sigma[*_{\mathcal{C}}\mathcal{C}]$.

(d) \Leftrightarrow (f) follows by 2.3.

(d) \Rightarrow (e) We show that for right \mathcal{C} -comodules M , any $*_{\mathcal{C}}$ -submodule N is a subcomodule. For this consider the map

$$\rho_N : N \rightarrow \text{Hom}_A(*_{\mathcal{C}}, N), \quad n \mapsto [f \mapsto f \cdot n].$$

With the inclusion $i : N \rightarrow M$, we have the commutative diagram with exact lines

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N & \xrightarrow{i} & M & \xrightarrow{p} & M/N & \longrightarrow & 0 \\ & & & & \downarrow \varrho_M & & & & \\ 0 & \longrightarrow & N \otimes_A \mathcal{C} & \xrightarrow{i \otimes I} & M \otimes_A \mathcal{C} & \xrightarrow{p \otimes I} & M/N \otimes_A \mathcal{C} & \longrightarrow & 0 \\ & & \downarrow \alpha_{N, \mathcal{C}} & & \downarrow \alpha_{M, \mathcal{C}} & & \downarrow \alpha_{M/N, \mathcal{C}} & & \\ 0 & \longrightarrow & \text{Hom}_A(*_{\mathcal{C}}, N) & \xrightarrow{\text{Hom}(*_{\mathcal{C}}, i)} & \text{Hom}_A(*_{\mathcal{C}}, M) & \longrightarrow & \text{Hom}_A(*_{\mathcal{C}}, M/N) & \longrightarrow & 0 \end{array},$$

where all the α 's are injective and $\text{Hom}(*_{\mathcal{C}}, i) \circ \rho_N = \alpha_{M, \mathcal{C}} \circ \varrho_M \circ i$. This implies $(p \otimes I) \circ \varrho_M \circ i = 0$, and by the kernel property, $\varrho_M \circ i$ factors through $N \rightarrow N \otimes_A \mathcal{C}$ thus yielding a \mathcal{C} -coaction on N .

The final assertion follows by 2.6 and the characterization of products in $\sigma[*_{\mathcal{C}}\mathcal{C}]$ (see 2.7). \square

As a corollary we can show when all $*_{\mathcal{C}}$ -modules are \mathcal{C} -comodules. This includes the reverse conclusion of [3, Lemma 4.3] and extends [11, Lemma 33].

3.6. $\mathcal{M}^{\mathcal{C}} = *_{\mathcal{C}}\mathcal{M}$.

For any A -coring \mathcal{C} , the following are equivalent:

(a) $\mathcal{M}^{\mathcal{C}} = *_{\mathcal{C}}\mathcal{M}$;

- (b) the functor $- \otimes_A \mathcal{C} : \mathcal{M}_A \rightarrow {}^*_{\mathcal{C}}\mathcal{M}$ has a left adjoint;
- (c) ${}_A\mathcal{C}$ is finitely generated and projective;
- (d) ${}_A\mathcal{C}$ is locally projective and \mathcal{C} is finitely generated as right \mathcal{C}^* -module.

Proof. (a) \Rightarrow (b) and (c) \Rightarrow (d) are obvious.

(b) \Rightarrow (c) By 2.6, $- \otimes_A \mathcal{C}$ preserves monomorphisms (injective morphisms) and hence ${}_A\mathcal{C}$ is flat. Moreover we obtain, for any family $\{M_\lambda\}_\Lambda$ in \mathcal{M}_A , the isomorphism

$$\left(\prod_{\Lambda} M_\lambda\right) \otimes_A \mathcal{C} \simeq \prod_{\Lambda} (M_\lambda \otimes_A \mathcal{C}),$$

which implies that ${}_A\mathcal{C}$ is finitely presented (e.g., [8, 12.9]) and hence projective.

(d) \Rightarrow (a) Recall that \mathcal{C}^* is the endomorphism ring of the faithful module ${}^*_{\mathcal{C}}\mathcal{C}$. Hence $\mathcal{C}_{\mathcal{C}^*}$ finitely generated implies $\mathcal{M}^{\mathcal{C}} = \sigma[{}^*_{\mathcal{C}}\mathcal{C}] = {}^*_{\mathcal{C}}\mathcal{M}$ (see 2.7). \square

Acknowledgement. The author is very grateful to Jawad Abuhlail for interesting and helpful discussions on the subject.

References

- [1] Abuhlail, J.Y., *Dualitätssätze für Hopf-Algebren über Ringen*, Dissertation, Universität Düsseldorf (2001)
- [2] Abuhlail, J.Y., Gómez-Torrecillas, J., Lobillo, F.J., *Duality and rational modules in Hopf algebras over commutative rings*, J. Algebra 240, 165-184 (2001)
- [3] Brzeziński, T., *The structure of corings*, Algebras and Repr. Theory, to appear
- [4] Garfinkel, G.S., *Universally torsionless and trace modules*, Trans. Amer. math. Soc. 215, 119-144 (1976)
- [5] Ohm, J., Bush, D.E., *Content modules and algebras*, Math. Scand. 31, 49-68 (1972)
- [6] Raynaud, M., Gruson, L., *Critère de platitude et de projectivité*, Inventiones Math. 13, 1-89 (1971)

- [7] Schubert, H., *Categories*, Springer, Berlin (1972)
- [8] Wisbauer, R., *Foundations of Module and Ring Theory*, Gordon and Breach, Reading, Paris (1991)
- [9] Wisbauer, R., *Semiperfect coalgebras over rings*, in *Algebras and Combinatorics, ICA'97*, Hong Kong, K.P. Shum, E. Taft, Z.X. Wan (ed), Springer Singapore, 487-512 (1999)
- [10] Wisbauer, R., *Weak Corings*, J. Algebra, to appear
- [11] Wischnewsky, M.B., *On linear representations of affine groups I*, Pac. J. Math. 61, 551-572 (1975)
- [12] Zimmermann-Huisgen, B., *Pure submodules of direct products of free modules*, Math. Ann. 224, 233-245 (1976)

Mathematisches Institut
Heinrich-Heine-Universität
40225 Düsseldorf
e-mail: wisbauer@math.uni-duesseldorf.de