

# Homologische Algebra

## Encyclopedia of Mathematics

[https://encyclopediaofmath.org/wiki/Homological\\_algebra](https://encyclopediaofmath.org/wiki/Homological_algebra)

Homological algebra – The branch of algebra whose main study is derived functors on various categories of algebraic objects (modules over a given ring, sheaves, etc.).

### AN INTRODUCTION TO HOMOLOGICAL ALGEBRA

JOSEPH J. ROTMAN  
Department of Mathematics  
University of Illinois  
Urbana, Illinois

Learning homological algebra is a two-stage affair. First, one must learn the language of  $\text{Ext}$  and  $\text{Tor}$  and what it describes. Second, one must be able to compute these things, and, often, this involves yet another language: spectral sequences. The following exercise appears on page 105 of Lang's book, "Algebra": "Take any book on homological algebra and prove all the theorems without looking at the proofs given in that book."

### AN INTRODUCTION TO HOMOLOGICAL ALGEBRA

CHARLES A. WEIBEL  
*Department of Mathematics  
Rutgers University*

Homological algebra is a tool used to prove nonconstructive existence theorems in algebra (and in algebraic topology). It also provides obstructions to carrying out various kinds of constructions; when the obstructions are zero, the construction is possible. Finally, it is detailed enough so that actual calculations may be performed in important cases. The following simple ques-

Bsp:  $G \curvearrowright A$

$$A^G := \{a \in A \mid g \cdot a = a\}$$

Frage:  $G \curvearrowright A, B$

$$A \twoheadrightarrow B$$

Ist dann auch  $A^G \twoheadrightarrow B^G$   
surjektiv?

Bsp:  $C_2 = \{\pm 1\}$   $C_2 \curvearrowright \mathbb{R}, \mathbb{Z}, \mathbb{Z}/4$

$$C_2 \curvearrowright S^1 \quad -1 \cdot \begin{array}{c} \updownarrow \end{array} \cdot 1$$

$$\begin{array}{ccc} \mathbb{Z}/15 & \twoheadrightarrow & \mathbb{Z}/3 \\ \cup & & \cup \\ \{0\} & \twoheadrightarrow & \{0\} \end{array}$$

✓

$$\begin{array}{ccc} \mathbb{Z} & \twoheadrightarrow & \mathbb{Z}/4 \\ \cup & & \cup \\ \{0\} & \twoheadrightarrow & \{0, 2\} \end{array}$$

✗

$$\begin{array}{ccc} \mathbb{R} & \twoheadrightarrow & \mathbb{R}/\mathbb{Z} \cong S^1 \\ \cup & & \cup \\ \{0\} & \twoheadrightarrow & \{0, \frac{1}{2}\} \end{array}$$

✗

Satz:  $A \xrightarrow{\pi} B$

Falls  $H^1(G; \ker \pi) = 0$ ,

dann

$$A^G \xrightarrow{\pi} B^G$$

## Bsp: spaltende Sequenzen

kurze exakte Sequenz abelscher Gruppen  $\sim A \hookrightarrow B \twoheadrightarrow B/A$

sie spaltet  $\sim B \cong A \oplus B/A$

Bsp:  $\mathbb{Z} \hookrightarrow \mathbb{Z} \oplus \mathbb{Z}/2 \twoheadrightarrow \mathbb{Z}/2$  spaltet  
 $x \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix} \mapsto x$

$2\mathbb{Z} \hookrightarrow \mathbb{Z} \twoheadrightarrow \mathbb{Z}/2$  spaltet nicht  
 $\parallel ?$

$\mathbb{Z} \oplus \mathbb{Z}/2 \leftarrow \exists x \neq 0: 2x = 0$

**Satz:** Genau dann spaltet jede exakte Sequenz

$$A \hookrightarrow ? \twoheadrightarrow C$$

wenn gilt:

$$\text{Ext}^1(C, A) = 0$$

Bsp (Weibell):

$B$  abelsche Gruppe  $n \in \mathbb{N}$   
 $\cup$   
 $A$

Wann ist  $nA = A \cap nB$ ?

Bsp:  $B = \mathbb{Z}$   $nB = 3\mathbb{Z}$   
 $\cup$   
 $A = 2\mathbb{Z}$   $nB \cap A = 3\mathbb{Z} \cap 2\mathbb{Z}$   
 $n = 3$   $= 6\mathbb{Z}$   
 $= nA$  ✓

$\overset{B/A}{\parallel}$   
 $\{b \in \mathbb{Z}/2 \mid 3b = 0\} = \{0\}$


Bsp:  $B = \mathbb{Z}$   $nB = 3\mathbb{Z}$   
 $\cup$   
 $A = 6\mathbb{Z}$   $nB \cap A = 6\mathbb{Z}$   
 $n = 3$   $nA = 18\mathbb{Z}$

$\overset{B/A}{\parallel}$   
 $\underbrace{\{b \in \mathbb{Z} \mid 3b = 0\}}_{\{0\}} \rightarrow \underbrace{\{b \in \mathbb{Z}/6 \mid 3b = 0\}}_{\{0, 2, 4\}}$

$$\text{Bsp: } B = S^1$$

$$A = C_4$$

$$n=3$$




$$nB = (S^1)^3 = S^1$$

$$nB \cap A = C_4$$

$$nA = (C_4)^3 = C_4$$

$$\begin{array}{ccc} \{b \in S^1 \mid b^3 = 1\} & \longrightarrow & \{b \in B/A \mid b^3 = 1\} \\ \parallel & & \parallel \\ \{1, \xi, \xi^2\} & \xrightarrow{\cong} & \{b \in S^1 \mid b^3 = 1\} = \{1, \xi, \xi^2\} \\ & \searrow & \\ & & \times 4 \end{array}$$



**Satz:**  $nA = A \cap nB$

genuau dann, wenn

$$\{b \in B \mid nb = 0\} \longrightarrow \{\bar{b} \in B/A \mid n\bar{b} = 0\}$$

surjektiv ist. Insbesondere ist

$$nA = A \cap nB$$

falls  $\{\bar{b} \in B/A \mid n\bar{b} = 0\} = 0$ .

[Beweis des Satzes:

①

$$\begin{array}{ccccc} uA \hookrightarrow A & \longrightarrow & A/uA \\ \downarrow & & \downarrow i & & \downarrow \bar{i} \\ uB \hookrightarrow B & \longrightarrow & B/uB \end{array}$$

$uA = uB \cap A \iff \bar{i}$  injektiv  
(Diagrammjagd)

②

$$A \hookrightarrow B \longrightarrow B/A$$

induziert

$$\text{Tor}(A, \mathbb{Z}/u) \longrightarrow \text{Tor}(B, \mathbb{Z}/u) \xrightarrow{p} \text{Tor}(B/A, \mathbb{Z}/u)$$

$$A/uA \xrightarrow{\bar{i}} B/uB \longrightarrow \frac{B/A}{u(B/A)}$$

lange exakte Sequenz

Daher:  $\bar{i}$  injektiv  $\iff p$  surjektiv


③

$$\text{Tor}(B, \mathbb{Z}/u) \xrightarrow{p} \text{Tor}(B/A, \mathbb{Z}/u)$$

$$\begin{array}{ccc} \parallel & & \parallel \\ \{b \in B \mid ub=0\} & & \{\bar{b} \in B/A \mid u\bar{b}=0\} \end{array} \quad ]$$



Meine Definition:

Homologische Algebra ist alles, was Sie brauchen, um meine Vorlesungen zur Algebraische Topologie (Topologie I & Topologie II) zu verstehen.



$$\text{Bsp: } S^1 \times S^1 \cong K \text{ ?}$$

Kleinsche Flasche




Tttrung, CC BY-SA 3.0,  
via Wikimedia Commons

Ansatz:

$$\begin{array}{ccc} S^1 \times S^1 & \xrightarrow{\quad} & H_i(S^1 \times S^1) \\ \text{~~HS~~} & & \\ K & \xrightarrow{\quad} & H_i(K) \end{array} \quad \begin{array}{l} \text{Homologie-} \\ \text{gruppen} \\ i = 0, 1, 2, \dots \end{array}$$

# Vorübung: $H_*(S^1)$



zellulärer Komplex

$$\mathbb{Z} \cdot a \oplus \mathbb{Z} \cdot b \oplus \mathbb{Z} \cdot c \xrightarrow{\delta} \mathbb{Z} \cdot A \oplus \mathbb{Z} \cdot B \oplus \mathbb{Z} \cdot C$$

$$\begin{array}{lcl} a & \mapsto & C - B \\ b & \mapsto & A - C \\ c & \mapsto & B - A \end{array}$$


$$\begin{array}{c} \mathbb{Z}^3 \xrightarrow{\delta} \mathbb{Z}^3 \\ \begin{pmatrix} 0 & -1 & -1 \\ -1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \\ \downarrow \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{array}$$

$$\begin{array}{c} H_1(S^1) \\ \parallel \\ \ker(\delta) \\ \parallel \wr \\ \mathbb{Z} \end{array}$$

$$\begin{array}{c} H_0(S^1) \\ \parallel \\ \mathbb{Z}^3 / \text{im } \delta \\ \parallel \\ \mathbb{Z} \end{array}$$



$$H_*(K)$$



zellulärer Komplex:

$$\begin{array}{c} 0 \\ \parallel \\ \mathbb{Z} \cdot C \oplus \mathbb{Z} \cdot D \xrightarrow{\delta_1} \mathbb{Z} \cdot a \oplus \mathbb{Z} \cdot b \oplus \mathbb{Z} \cdot c \xrightarrow{\delta_0} \mathbb{Z} \cdot A \end{array}$$

$$\begin{array}{ll} C \mapsto a+b-c & a \mapsto 0 \\ D \mapsto c+a-b & b \mapsto 0 \\ & c \mapsto 0 \end{array}$$

$$\mathbb{Z}^2 \xrightarrow{\begin{pmatrix} 1 & 1 \\ 1 & -1 \\ -1 & 1 \end{pmatrix}} \mathbb{Z}^3 \xrightarrow{0} \mathbb{Z}$$

$$\vdots$$

$$\begin{pmatrix} 2 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\begin{array}{c} H_2(K) \\ \parallel \\ \ker(\delta_1) \\ \parallel \\ 0 \end{array}$$

$$\begin{array}{c} H_1 \\ \parallel \\ \ker(\delta_0) \\ \hline \text{im}(\delta_1) \\ \parallel \\ \mathbb{Z}^3 \\ \hline 2\mathbb{Z} \oplus \mathbb{Z} \oplus 0 \\ \parallel \\ \mathbb{Z}/2 \oplus \mathbb{Z} \end{array}$$

$$\begin{array}{c} H_0(K) \\ \parallel \\ \mathbb{Z} / \text{im} \delta_0 \\ \parallel \\ \mathbb{Z} \end{array}$$

$$H_*(S^1 \times S^1)$$

## Algebraic Topology

Allen Hatcher

top. spaces

**Theorem 3B.6.** If  $X$  and  $Y$  are ~~CW complexes~~ and  $R$  is a ~~principal ideal domain~~, then there are natural short exact sequences

$$0 \rightarrow \bigoplus_i (H_i(X; R) \otimes_R H_{n-i}(Y; R)) \rightarrow H_n(X \times Y; R) \rightarrow \bigoplus_i \text{Tor}_R(H_i(X; R), H_{n-i-1}(Y; R)) \rightarrow 0$$

and these sequences split.

In unserem Fall  $X = Y = S^1$  ist  
jeweils  $\text{Tor}(\dots, \dots) = 0$  und wir  
erhalten

$$H_*(S^1 \times S^1) = H_*(S^1) \oplus H_*(S^1)$$

$$= \begin{pmatrix} \mathbb{Z} \\ \mathbb{Z} \end{pmatrix} \oplus \begin{pmatrix} \mathbb{Z} \\ \mathbb{Z} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} \end{pmatrix} \begin{matrix} \mathbb{Z} = 1+1 \\ 1 = 1+0 = 0+1 \\ 0 = 0+0 \end{matrix}$$

$$(\mathbb{Z} \oplus \mathbb{Z} = \mathbb{Z})$$

$$H_2(S^1 \times S^1)$$

$\parallel$

$$\mathbb{Z}$$

$$H_1(S^1 \times S^1)$$

$\parallel$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$H_0(S^1 \times S^1)$$

$\parallel$

$$\mathbb{Z}$$

Offenbar

$$H_2(K) \neq H_2(S^1 \times S^1)$$

(und

$$H_1(K) \neq H_1(S^1 \times S^1))$$

also

$$K \neq S^1 \times S^1 \text{ (nicht homöomorph)}$$