

Homologische Algebra

Encyclopedia of Mathematics

https://encyclopediaofmath.org/wiki/Homological_algebra

Homological algebra – The branch of algebra whose main study is derived functors on various categories of algebraic objects (modules over a given ring, sheaves, etc.).

AN INTRODUCTION TO HOMOLOGICAL ALGEBRA

JOSEPH J. ROTMAN
Department of Mathematics
University of Illinois
Urbana, Illinois

Learning homological algebra is a two-stage affair. First, one must learn the language of Ext and Tor and what it describes. Second, one must be able to compute these things, and, often, this involves yet another language: spectral sequences. The following exercise appears on page 105 of Lang's book, "Algebra": "Take any book on homological algebra and prove all the theorems without looking at the proofs given in that book."

AN INTRODUCTION TO HOMOLOGICAL ALGEBRA

CHARLES A. WEIBEL
*Department of Mathematics
Rutgers University*

Homological algebra is a tool used to prove nonconstructive existence theorems in algebra (and in algebraic topology). It also provides obstructions to carrying out various kinds of constructions; when the obstructions are zero, the construction is possible. Finally, it is detailed enough so that actual calculations may be performed in important cases. The following simple ques-

Bsp: $G \curvearrowright A$

$$A^G := \{a \in A \mid g \cdot a = a\}$$

Frage: $G \curvearrowright A, B \quad A \twoheadrightarrow B$

Ist dann auch $A^G \twoheadrightarrow B^G$
surjektiv?

Bsp: $C_2 = \{\pm 1\} \quad C_2 \curvearrowright \mathbb{R}, \mathbb{Z}, \mathbb{Z}/4$

$$C_2 \curvearrowright S^1 \quad -1 \begin{pmatrix} \updownarrow \end{pmatrix} 1$$

$$\begin{array}{ccc} \mathbb{Z}/15 & \twoheadrightarrow & \mathbb{Z}/3 \\ \cup & & \cup \\ \{0\} & \twoheadrightarrow & \{0\} \end{array}$$

✓

$$\begin{array}{ccc} \mathbb{Z} & \twoheadrightarrow & \mathbb{Z}/4 \\ \cup & & \cup \\ \{0\} & \twoheadrightarrow & \{0, 2\} \end{array}$$

✗

$$\begin{array}{ccc} \mathbb{R} & \twoheadrightarrow & \mathbb{R}/\mathbb{Z} \cong S^1 \\ \cup & & \cup \\ \{0\} & \twoheadrightarrow & \{0, \frac{1}{2}\} \end{array} \quad \begin{array}{ccc} & & \cup \\ & & \{\pm 1\} \end{array}$$

✗

Satz: $A \xrightarrow{\pi} B$

Falls $H^1(G; \ker \pi) = 0$,

dann

$$A^G \xrightarrow{\pi} B^G$$

Bsp: spaltende Sequenzen

kurze exakte Sequenz abelscher
Gruppen $\sim A \hookrightarrow B \twoheadrightarrow B/A$

sie spaltet $\sim B \cong A \oplus B/A$

Bsp: $\mathbb{Z} \hookrightarrow \mathbb{Z} \oplus \mathbb{Z}/2 \twoheadrightarrow \mathbb{Z}/2$ spaltet
 $x \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix} \mapsto x$

$2\mathbb{Z} \hookrightarrow \mathbb{Z} \twoheadrightarrow \mathbb{Z}/2$ spaltet nicht
 $\parallel ?$

$\mathbb{Z} \oplus \mathbb{Z}/2 \leftarrow \exists x \neq 0: 2x = 0$

Satz: Genau dann spaltet
jede exakte Sequenz

$$A \hookrightarrow ? \twoheadrightarrow C$$

wenn gilt:

$$\text{Ext}^1(C, A) = 0$$

Bsp (Weibel):

B abelsche Gruppe $u \in \mathbb{N}$
 \cup
 A

Wann ist $uA = A \cap uB$?


Bsp: $B = \mathbb{Z}$ $uB = 3\mathbb{Z}$
 \cup
 $A = 2\mathbb{Z}$ $uB \cap A = 3\mathbb{Z} \cap 2\mathbb{Z}$
 $u = 3$ $= 6\mathbb{Z}$
 $= uA$ ✓

$$\begin{array}{c} B/A \\ \parallel \\ \{b \in \mathbb{Z}/2 \mid 3b = 0\} = \{0\} \end{array}$$


Bsp: $B = \mathbb{Z}$ $uB = 3\mathbb{Z}$
 \cup
 $A = 6\mathbb{Z}$ $uB \cap A = 6\mathbb{Z}$
 $u = 3$ $uA = 18\mathbb{Z}$

$$\begin{array}{c} B/A \\ \parallel \\ \underbrace{\{b \in \mathbb{Z} \mid 3b = 0\}}_{\{0\}} \rightarrow \underbrace{\{b \in \mathbb{Z}/6 \mid 3b = 0\}}_{\{0, 2, 4\}} \end{array}$$

Bsp: $B = S^1$ $\cup B = (S^1)^3 = S^1$
 $A = C_4$ $\cup B \cap A = C_4$
 $n=3$ $\cup A = (C_4)^3 = C_4$



$\{b \in S^1 \mid b^3 = 1\} \rightarrow \{b \in B/A \mid b^3 = 1\}$
 $\{1, \xi, \xi^2\}$ \cong $\{b \in S^1 \mid b^3 = 1\} = \{1, \xi, \xi^2\}$
 \times $\xrightarrow{\cong}$ $\times 4$



Satz: $\cup A = A \cap \cup B$

genau dann, wenn

$\{b \in B \mid \cup b = 0\} \rightarrow \{\bar{b} \in B/A \mid \cup \bar{b} = 0\}$

surjektiv ist. Insbesondere ist

$\cup A = A \cap \cup B$

falls $\{\bar{b} \in B/A \mid \cup \bar{b} = 0\} = 0$.

[Beweis des Satzes:

$$\textcircled{1} \quad \begin{array}{ccccc} uA \hookrightarrow A & \longrightarrow & A/uA \\ \downarrow & & \downarrow \bar{i} \\ uB \hookrightarrow B & \longrightarrow & B/uB \end{array}$$

$uA = uB \cap A \iff \bar{i}$ injektiv
(Diagrammjagd)

$$\textcircled{2} \quad A \hookrightarrow B \longrightarrow \frac{B}{A}$$

induziert

$$\text{Tor}(A, \frac{\mathbb{Z}}{n}) \longrightarrow \text{Tor}(B, \frac{\mathbb{Z}}{n}) \xrightarrow{P} \text{Tor}(\frac{B}{A}, \frac{\mathbb{Z}}{n})$$

$$\frac{A}{uA} \xrightarrow{\bar{i}} \frac{B}{uB} \longrightarrow \frac{B/A}{u(B/A)}$$

lange exakte Sequenz


Daher: \bar{i} injektiv \iff P surjektiv

$$\textcircled{3} \quad \text{Tor}(B, \frac{\mathbb{Z}}{n}) \xrightarrow{P} \text{Tor}(\frac{B}{A}, \frac{\mathbb{Z}}{n})$$



$$\begin{array}{ccc} \parallel & & \parallel \\ \{b \in B \mid ub=0\} & & \{\bar{b} \in B/A \mid u\bar{b}=0\} \end{array}]$$

Meine Definition:

Homologische Algebra ist alles, was Sie brauchen, um meine Vorlesungen zur Algebraische Topologie (Topologie I & Topologie II) zu verstehen.



Bsp: $S^1 \times S^1 \cong K$?
Kleinsche Flasche




Tttrung, CC BY-SA 3.0, via Wikimedia Commons

Ansatz:

$$\begin{array}{ccc} S^1 \times S^1 & \xrightarrow{\quad} & H_i(S^1 \times S^1) \\ \text{Hs} & & \text{Hs} \\ K & \xrightarrow{\quad} & H_i(K) \end{array} \quad \begin{array}{l} \text{Homologie-} \\ \text{gruppen} \\ i = 0, 1, 2, \dots \end{array}$$

Vorübung: $H_*(S^1)$



zellulärer Komplex

$$\mathbb{Z} \cdot a \oplus \mathbb{Z} \cdot b \oplus \mathbb{Z} \cdot c \xrightarrow{\delta} \mathbb{Z} \cdot A \oplus \mathbb{Z} \cdot B \oplus \mathbb{Z} \cdot C$$

$$a \mapsto C - B$$

$$b \mapsto A - C$$

$$c \mapsto B - A$$

$$\mathbb{Z}^3 \xrightarrow{\delta} \mathbb{Z}^3$$

$$\begin{pmatrix} 0 & -1 & -1 \\ -1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$$

δ

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$H_1(S^1)$$

$$\cong \ker(\delta)$$

$$\cong \mathbb{Z}$$


$$H_0(S^1)$$

$$\cong \mathbb{Z}^3 / \text{im } \delta$$

$$\cong$$

$$\mathbb{Z}$$

$H_*(K)$



zellulärer Komplex:

$$\mathbb{Z} \cdot C \oplus \mathbb{Z} \cdot D \xrightarrow{\delta_1} \mathbb{Z} \cdot a \oplus \mathbb{Z} \cdot b \oplus \mathbb{Z} \cdot c \xrightarrow{\delta_0} \mathbb{Z} \cdot A$$

$$\begin{array}{l} C \mapsto a+b-c \quad a \mapsto 0 \\ D \mapsto c+a-b \quad b \mapsto 0 \\ \quad \quad \quad \quad \quad c \mapsto 0 \end{array}$$

$$\mathbb{Z}^2 \xrightarrow{\begin{pmatrix} 1 & 1 \\ 1 & -1 \\ -1 & 1 \end{pmatrix}} \mathbb{Z}^3 \xrightarrow{0} \mathbb{Z}$$

⋮

$$\begin{pmatrix} 2 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\begin{array}{l} H_2(K) \\ \parallel \\ \ker(\delta_1) \\ \parallel \\ 0 \end{array}$$

$$\begin{array}{l} H_1 \\ \parallel \\ \ker(\delta_0) \\ \hline \text{im}(\delta_1) \\ \parallel \\ \mathbb{Z}^3 \\ \hline 2\mathbb{Z} \oplus \mathbb{Z} \oplus 0 \\ \parallel \\ \mathbb{Z}/2 \oplus \mathbb{Z} \end{array}$$

$$\begin{array}{l} H_0(K) \\ \parallel \\ \mathbb{Z} / \text{im} \delta_0 \\ \parallel \\ \mathbb{Z} \end{array}$$

$$H_* (S^1 \times S^1)$$

Algebraic Topology

Allen Hatcher

top. spaces

Theorem 3B.6. If X and Y are CW-complexes and R is a principal ideal domain, then there are natural short exact sequences

$$0 \rightarrow \bigoplus_i (H_i(X; R) \otimes_R H_{n-i}(Y; R)) \rightarrow H_n(X \times Y; R) \rightarrow \bigoplus_i \text{Tor}_R(H_i(X; R), H_{n-i-1}(Y; R)) \rightarrow 0$$

and these sequences split.

In unserem Fall $X = Y = S^1$ ist jeweils $\text{Tor}(\dots, \dots) = 0$ und wir erhalten

$$H_* (S^1 \times S^1) = H_* (S^1) \oplus H_* (S^1)$$

$$= \begin{pmatrix} \mathbb{Z} \\ \mathbb{Z} \end{pmatrix} \oplus \begin{pmatrix} \mathbb{Z} \\ \mathbb{Z} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} \end{pmatrix} \begin{matrix} 2 = 1+1 \\ 1 = 1+0 = 0+1 \\ 0 = 0+0 \end{matrix}$$

$$(\mathbb{Z} \oplus \mathbb{Z} = \mathbb{Z})$$

$$H_2(S^1 \times S^1)$$

\cong

$$\mathbb{Z}$$

$$H_1(S^1 \times S^1)$$

\cong

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$H_0(S^1 \times S^1)$$

\cong

$$\mathbb{Z}$$

Offenbar $H_2(K) \neq H_2(S^1 \times S^1)$

(und $H_1(K) \neq H_1(S^1 \times S^1)$)

also $K \neq S^1 \times S^1$ (nicht
homöomorph)